PSAP Pharmacotherapy Self-Assessment Program

REFERENCE VALUES FOR COMMON LABORATORY TESTS¹

| Serum Chemistries | Reference Range | | |
|---|--------------------------------------|--|--|
| Alanine aminotransferase (ALT) | 10–40 U/L | | |
| Albumin | 3.5–5 g/dL (adults) | | |
| | 3.4–4.2 g/dL (young children) | | |
| Alkaline phosphatase (ALP) | Varies with age: | | |
| | 30–120 IU/L (adults) | | |
| | 150–420 IU/L (children) | | |
| Ammonia | 15–45 mcg/dL | | |
| Amylase | 27–131 U/L | | |
| Aspartate aminotransferase (AST) | 10–30 U/L | | |
| Bilirubin, direct | 0.1–0.3 mg/dL | | |
| Bilirubin, total | 0.3–1.2 mg/dL | | |
| Blood urea nitrogen (BUN) | 8–23 mg/dL (adults) | | |
| | Lower in children | | |
| Calcium, ionized | 4.6–5.1 mg/dL | | |
| Calcium, total serum | 8.2–10.2 mg/dL | | |
| Carbon dioxide (venous) (CO ₂) | 22–28 mEq/L | | |
| Chloride (Cl) | 96–106 mEq/L | | |
| C-reactive protein (CRP) | 0.08–3.1 mg/L | | |
| Creatinine kinase (CK) | 40–150 U/L | | |
| Creatinine, serum (SCr) | 0.6–1.2 mg/dL (adults) | | |
| | 0.2–0.7 mg/dL (children) | | |
| Creatinine (clearance) (CrCl) | 75–125 mL/minute/1.73 m ² | | |
| Ferritin | 15–200 ng/mL | | |
| γ-Glutamyl transpeptidase | 2–30 U/L | | |
| Glucose, serum | 70–110 mg/dL | | |
| Hemoglobin A1C percent of total hemoglobin | 4%-7% | | |
| Lactate dehydrogenase (LDH) | 100–200 U/L | | |
| Lipase | 31–186 U/L | | |
| Magnesium | 1.3–2.1 mEq/L | | |
| Osmolality, serum | 275–295 mOsm/kg | | |
| Phosphorus | 2.3–4.7 mg/dL (adults) | | |
| - | 3.7–5.6 mg/dL (children) | | |
| Potassium | 3.5–5.0 mEq/L | | |
| Prealbumin | 19.5–35.8 mg/dL | | |
| Sodium | 136–142 mEg/L | | |
| Uric acid, serum | 4–8 mg/dL | | |
| • | | | |

¹Values given in this table are commonly accepted reference ranges compiled from several sources. Patient-specific goals may differ depending on age, sex, clinical condition, and the laboratory methodology used to perform the assay.

Information from: AMA Manual of Style. Available at www.Amamanualofstyle.com/page/si-conversion-calculator. Last accessed 11/21/2018); Lee M, ed. Basic Skills in Interpreting Laboratory Data, 6th ed. Bethesda, MD: American Society of Health-System Pharmacists, 2017; and DiPiro JT, Talbert RL, Yee GC, et al., eds. Pharmacotherapy: A Pathophysiologic Approach, 10th ed. New York: McGraw-Hill, 2017.

| Hematology/Coagulation | Reference Range | | |
|---|---|--|--|
| Hematocrit (Hct) | 42%–50% (men) | | |
| | 36%–45% (women) | | |
| Hemoglobin (Hgb) | 14–18 g/dL (men) | | |
| | 12–16 g/dL (women) | | |
| International normalized ratio (INR) | 0.9–1.1 | | |
| Mean corpuscular hemoglobin (MCH) | 26–34 pg/cell | | |
| Mean corpuscular hemoglobin concentration (MCHC) | 33–37 g/dL | | |
| Mean corpuscular volume (MCV) | 80–100 fL/cell | | |
| Partial thromboplastin time (PTT) | 25–40 seconds | | |
| Platelet count (Plt) | 150,000–350,000 cells/mm ³ | | |
| Prothrombin time (PT) | 10–13 seconds | | |
| Red blood cell count (RBC) | 4.5–5.9 × 10 ⁶ cells/mm ³ (men) | | |
| | 4.1–5.1 × 10 ⁶ cells/mm ³ (women) | | |
| Reticulocyte percent of red blood cells | 0.5%-1.5% | | |
| White blood cell count (WBC) | 4.5–11.0 × 10 ³ cells/mm ³ | | |
| | | | |

| Comme Linida | Deference Dence | |
|---|-------------------|-------------|
| Serum Lipids | Reference Range | |
| Cholesterol, total (TC), | < 200 mg/dL | |
| desirable | | |
| High-density lipoprotein (HDL) | <u>≥</u> 60 mg/dL | |
| cholesterol, desirable | | |
| Low-density lipoprotein (LDL) | < 100 mg/dL | |
| cholesterol | | |
| Triglycerides (TG) | < 150 mg/dL | |
| Blood Gases | Arterial | Venous |
| рН | 7.35–7.45 | 7.31-7.41 |
| Partial pressure of carbon | 35–45 mm Hg | 40–52 mm Hg |
| dioxide (PCO ₂) | | |
| Partial pressure of oxygen (PO ₂) | 80–100 mm Hg | 30–50 mm Hg |
| Oxygen saturation (SaO ₂) | > 90% | 60%-75% |
| Serum bicarbonate (HCO ₃) | 22–26 mEq/L | 21–28 mEq/L |
| Urinalysis | | |
| Leukocyte esterase, nitrite, | Negative | |
| protein, blood, ketones, | | |
| bilirubin, glucose | | |
| рН | 4.5-8.0 | |
| Specific gravity | 1.010–1.025 | |

